

Current Carrying States in a Random Magnetic Field

Kun Yang and R. N. Bhatt

Department of Electrical Engineering, Princeton University, Princeton, NJ 08544

(February 1, 2008)

We report results of a numerical study of noninteracting electrons moving in two dimensions, in the presence of a random potential and a random magnetic field for a sequence of finite sizes, using topological properties of the wave functions to identify extended states. Our results are consistent with the existence of a second order localization-delocalization transition driven by the random potential. The critical randomness strength and localization length exponent are estimated via a finite size scaling analysis.

71.30.+h, 73.40.Hm

The problem of Anderson localization in two dimensions (2D) continues to be the subject of extensive research recently. Scaling arguments¹ predict that all one-particle states are localized in 2D, in the presence of scattering potential which respects time reversal symmetry. In the presence of a *uniform* magnetic field, however, because of the fact that the magnetic field breaks time reversal symmetry and suppresses the enhanced back scattering, extended states are found at individual critical energies.² These extended states are crucial to the existence of the integer quantum Hall effect.²

Recently there has been considerable interest in the localization problem in 2D in the presence of a *random* magnetic field with *zero* mean, motivated by the work of Halperin, Lee and Read,³ and Kalmeyer and Zhang,⁴ on the problem of the half filled Landau level. These authors argue that if one treats the electrons as composite fermions⁵ carrying two flux quanta, the flux carried by the composite fermions cancels that of the external magnetic field *on average* when the Landau level filling factor is 1/2, and the systems may be viewed as a Fermi liquid like metal. Disorder induces local fluctuations in the density of the composite fermion, and hence at the mean field level the composite fermions see a *fluctuating* local magnetic field with zero mean, in the presence of disorder.

Following this development, there have been many efforts devoted to the study of the localization problem of noninteracting particles moving in a random magnetic field with zero mean.^{6–16} However, the key issue, namely whether extended states exist in the thermodynamic limit, remains unresolved. Zhang and Arovas¹⁰ argued that in the field theory description of the problem, the local magnetic flux gives rise to a local topological density which has long range interaction and induces a localization-delocalization transition (see, however, Ref. [11]). Their conclusion is supported by numerical work of several groups^{7,8,12,13} which found extended states near the band center on lattice models with weak random potential and finite random magnetic field, although Sugiyama and Nagaosa⁶ concluded otherwise. On the other hand, Lee and Chalker⁹ simulated a network model which they argue to be appropriate to the problem of random magnetic field, and find states are lo-

calized at all energies. Subsequently Kim, Furusaki and Lee¹⁴ showed that the network model of Lee and Chalker may be mapped onto an $SU(2N)$ spin chain in the limit $N \rightarrow 0$, which has short range correlations and an excitation gap; they thus confirm the conclusion of Lee and Chalker within the framework of the network model. The network model was originally introduced¹⁷ in the study of the integer quantum Hall effect, and has enjoyed tremendous success in that context. Such success is ensured by the fact that in the presence of a strong *uniform* magnetic field, the magnetic length is *very small*, thus one body states live along *one dimensional* equipotential lines. In the case of a *random* magnetic field, however, the relevant percolating equal field lines are along $B = 0$, where the magnetic length is *infinite*! Therefore it is not completely obvious that the network model of Lee and Chalker is an appropriate description of the random magnetic field problem.

In this paper we report a systematic finite size numerical study of the electron localization problem in the presence of a random magnetic field with zero mean, as well as a random potential, on the square lattice. We study topological properties of one-electron wave functions by calculating their topological quantum numbers called the Chern number.^{18,19} This approach has proved very successful in the study of integer quantum Hall effect and transitions.^{20–22} In particular, finite size corrections appear to be much less in this approach, compared with others, implying a rapid convergence to the thermodynamic limit. Recently it has been applied by Sheng and Weng¹³ to the present problem. There are, however, several important differences between our work and theirs. Firstly, their work as well as other numerical studies,^{6–8,12} concentrate on systems with a relatively *weak* (or zero) random potential, and attempt to identify a mobility edge. When the strength of random potential is weak, the length scales involved may be quite large, and it is often hard to distinguish whether the localization length near the band center is truly infinite or very large but finite. In contrast, we study systems with a series of different random potential strengths, with the same strength of random *flux*. The idea is that *if* extended states do exist for weak randomness, there must exist a critical random potential strength (which is typically *not*

very weak) at which all extended states disappear, and scaling behavior should be observable near this critical point. Secondly, unlike Sheng and Weng, who study behavior of density of states ($\rho(E)$) and density of current carrying states ($\rho_c(E)$), which are functions of energy E , we focus on quantities like the *total number* of extended states N_c in a finite size system. N_c is the zeroth moment of the distribution function $\rho_c(E)$. In other random systems such as spin glasses low order moments of distribution functions are known to converge to an acceptable level of accuracy relatively fast with sample averaging; the higher moments and the distributions themselves are still quite noisy at that stage. Finally, our massively parallel computer allows us to study samples of considerably larger sizes than those in Ref. [13].

We consider the one body tight binding Hamiltonian on the square lattice:

$$H = - \sum_{\langle ij \rangle} (e^{ia_{ij}} c_i^\dagger c_j + e^{-ia_{ij}} c_j^\dagger c_i) + \sum_i \epsilon_i c_i^\dagger c_i, \quad (1)$$

where c_i is the fermion operator on lattice site i . The first term represents hopping or kinetic energy of the electrons, and the summation is over nearest neighbors. The flux through each square is equal to the summation of a_{ij} along its four edges. We take the flux to be random and uniformly distributed from $-h\pi$ and $h\pi$, where $0 \leq h \leq 1$. The second term represents a random onsite potential. We take ϵ_i to be uniformly distributed between $-w$ and w . h and w parameterize the strength of random magnetic field and random potential respectively; when $h = 1$ the flux through each square is completely random. We have set the hopping matrix element to be 1 and use it as the unit of energy. We study samples of square geometry with linear size L (with number of sites $N_s = L^2$), and impose periodic boundary conditions in both directions: $\Psi(k + L\hat{x}) = e^{i\phi_1} \Psi(k)$, and $\Psi(k + L\hat{y}) = e^{i\phi_2} \Psi(k)$. The Hall conductance of an individual eigenstate $|m\rangle$ can be obtained easily using the Kubo formula:

$$\sigma_{xy}^m = \frac{ie^2\hbar}{N_s} \sum_{n \neq m} \frac{\langle m | v_y | n \rangle \langle n | v_x | m \rangle - \langle m | v_x | n \rangle \langle n | v_y | m \rangle}{(E_n - E_m)^2}, \quad (2)$$

where E_n is the energy of the n th state, and

$$v_\tau = (i/\hbar) \sum_j (c_{j+\tau}^\dagger c_j e^{ia_{j+\tau,j}} - c_j^\dagger c_{j+\tau} e^{-ia_{j+\tau,j}}) \quad (3)$$

is the velocity operator along direction τ ($= \hat{x}$ or $\hat{y}$). σ_{xy} is identically zero in the absence of magnetic flux, as guaranteed by time reversal symmetry. In the presence of random magnetic flux with zero average, the time reversal symmetry is broken in a specific configuration of randomness, and *individual* states may have nonzero σ_{xy} , although the *disorder averaged* Hall conductance is always zero for any given Fermi energy, since the averaging process restores the time reversal symmetry. σ_{xy}^m

depends on the boundary conditions imposed. As shown by Niu *et al.*, the boundary condition averaged Hall conductance takes the form¹⁹

$$\langle \sigma_{xy}^m \rangle = \frac{1}{4\pi^2} \int d\phi_1 d\phi_2 \sigma_{xy}^m(\phi_1, \phi_2) = C(m)e^2/h, \quad (4)$$

where $C(m)$ is an integer called the Chern number of the state $|m\rangle$. States with nonzero Chern numbers carry Hall current and are necessarily extended states.²⁰ Thus by numerically diagonalizing the Hamiltonian on a grid of ϕ_1 and ϕ_2 , and calculating the Chern numbers of states of finite size systems by converting the integral in (2) to a sum over grid points, we are able to identify extended states in a *finite size* system unambiguously.

In this work we have studied finite size samples with linear size ranging from $L = 4$ to $L = 19$, for different random potential and random field strength. For each particular randomness strength and system size, we average over 30 to 2000 different configurations of randomness. Depending on system size and randomness strength, the number of grids necessary to determine the Chern numbers unambiguously using Eq. (4) ranges from 35×35 to 70×70 .

We plot in Fig. 1 the dependence of disorder averaged number of current carrying states N_c versus the system size N_s , for $h = 0.5$ and a series of different w , on a *double logarithmic* plot. We find for a given system size, N_c decreases as w increases. This is exactly what is expected, since random potential tends to localize states. For very strong randomness ($w \geq 4.5$) and big enough system size, we see N_c decreases with increasing N_s , indicating there are no extended states in the thermodynamic limit, and the localization length is finite throughout the band. For somewhat weaker randomness, *e.g.* $w = 4.0$, although N_c increases with N_s within the size range we are able to reach, the curves obviously have downward curvature, and beyond certain N_s , N_c presumably decreases with increasing size and eventually goes to zero. Current carrying states appear in *finite size* systems in this case because the localization length ξ for states near the band center, although finite, may be larger than the system size L .²¹ For even weaker randomness ($w < 3.0$), however, we do not see any evidence of such downward curvature, and it appears that $N_c \rightarrow \infty$ as $N_s \rightarrow \infty$. Such behavior is expected if extended states exist in the thermodynamic limit, either at individual critical energies,^{21,22} or in a finite width band near the band center. In both cases, one expects the asymptotic behavior on the double logarithmic plot to be linear. In the former case, the slope is expected to be less than 1, while for a band of extended states we have $N_c \propto N_s$, hence the slope equals unity. For weak enough random potential, $w \leq 2.0$, we do find that the slope y is extremely close to 1, consistent with a band of extended states in the thermodynamic limit.

It has been suggested¹¹ that for weak randomness states are just barely localized (as in the case without magnetic field¹) with exponentially large ξ near the band

center, much larger than system sizes accessible to numerical studies. While this may give the impression of an extended band, one would expect a slow, logarithmic decrease in N_c/N_s with increasing system size. However, we find no such evidence for small w ; N_c/N_s appears to be a constant, or *increases* slightly. If, on the other hand, there is a band of extended states for small w , the width of such a band should decrease as w increases and shrink to zero at a critical random potential w_c , above which ξ is finite throughout the band (see Fig. 2). For $w < w_c$ we should have $N_c \propto N_s$ as $N_s \rightarrow \infty$, while for $w > w_c$, $N_c \rightarrow 0$ in this limit. Right at the critical point $w = w_c$, there is a single critical energy at which the localization length diverges, just like the case of a uniform magnetic field,^{21,22} and we should have $N_c \sim N_s^{y_c}$, where y_c is an exponent related to the localization length exponent ν through^{21–23}

$$y_c = 1 - \frac{1}{4\nu}. \quad (5)$$

For w close to w_c , there is a characteristic length scale that scales as $\xi_m \sim |w - w_c|^{-\nu}$. For $w > w_c$, ξ_m is the maximum localization length of the system, while for $w < w_c$, ξ_m is the length scale above which the behavior $N_c \propto N_s$ is seen. Near the critical point we should see scaling behavior:

$$N_c \sim N_s^{y_c} \tilde{F}^{\pm}(L/\xi_m) \sim N_s^{y_c} F^{\pm}(N_s|w - w_c|^{2\nu}), \quad (6)$$

where $F^{\pm}(x)$ are two different scaling functions for $w < w_c$ and $w > w_c$ respectively, which should satisfy the asymptotic behavior $F^+(x) \sim x^{1-y_c}$ and $F^-(x) \rightarrow 0$ as $x \rightarrow \infty$. Therefore, *if* there is indeed a critical randomness w_c , we ought to see scaling behavior (6), by tuning a *single* parameter w_c . The other parameter ν in the scaling relation (6) may be determined using Eq. (5), since the slope y is known from the data presented in Fig. 1.

Such scaling behavior is indeed seen, and the data collapses best for $w_c = 3.0$ and $\nu = 1.8$, as shown in Fig. 3. At $w = 3.0$ the linear fit of $\log N_c$ versus $\log N_s$ yields $y = 0.87 \pm 0.01$, and from Eq. (5) we obtain $\nu = 1.9 \pm 0.2$, consistent with scaling results. We emphasize that data collapsing of *two* different scaling curves is achieved by tuning *one* parameter, due to the constraint Eq. (5). The scaling behavior we see here supports the existence of a band of extended states for weak random potential, and would not be expected if there were no localization-delocalization transition at finite w_c . In order to make sure that our data are indeed in the scaling regime, we plot in Fig. 4 both the total density of states $\rho(E)$ and density of current carrying states $\rho_c(E)$ (per site), for $w = 3.0$ and $h = 0.5$, for two different system sizes $L = 4$ and $L = 16$. We find the width of $\rho_c(E)$ is considerably smaller than that of $\rho(E)$, especially for the larger size. Further the width of $\rho_c(E)$ is size dependent while that of $\rho(E)$ is essentially the same for both sizes. This is similar to what is seen in the study of the localization problem in lowest Landau level,²¹ and indicates that we are indeed

in the scaling regime. By contrast the width of $\rho_c(E)$ and $\rho(E)$ is almost the same for Gaussian white noise potential in the first Landau level, where the localization lengths are known to be very large.^{2,24}

From the scaling we are also able to estimate the critical randomness to be $w_c = 3.0 \pm 0.2$, and $\nu = 1.8 \pm 0.2$, for $h = 0.5$. The localization length exponent ν we obtain here is very close to that estimated by Sheng and Weng¹³ using a very different scaling scheme, although they do not have a quantitative estimate of the critical random potential strength w_c .

In summary, we have found good evidence supporting the existence of a localization-delocalization transition driven by random potential, in the 2D square lattice in the presence of a random magnetic field with zero average, implying the existence of extended states at weak randomness. The critical random potential strength and localization length exponent are estimated using finite size scaling analysis.

We thank Yong Baek Kim, Z. Y. Weng, and P. Wölfle for helpful discussions. This work was supported by NSF grants DMR-9400262 and CDA-9121709. R.N.B. was also supported in part by a Guggenheim fellowship, and thanks the Aspen Center for Physics for hospitality while this manuscript was being written up.

-
- ¹ E. Abrahams, P. W. Anderson, D. C. Licciardello and T. V. Ramakrishnan, Phys. Rev. Lett. **42**, 673 (1979).
 - ² B. Huckestein, Rev. Mod. Phys. **67**, 357 (1995).
 - ³ B. I. Halperin, P. A. Lee and N. Read, Phys. Rev. B **47**, 7312 (1993).
 - ⁴ V. Kalmeyer and S. C. Zhang, Phys. Rev. B **46**, 9889 (1992).
 - ⁵ J. K. Jain, Phys. Rev. Lett. **63**, 199 (1989).
 - ⁶ T. Sugiyama and N. Nagaosa, Phys. Rev. Lett. **70**, 1980 (1993).
 - ⁷ Y. Avishai, Y. Hatsugai, and M. Kohmoto, Phys. Rev. B **47**, 9561 (1993).
 - ⁸ V. Kalmeyer, D. Wei, D. P. Arovas, and S. C. Zhang, Phys. Rev. B **48**, 11095 (1993).
 - ⁹ D. K. K. Lee and J. T. Chalker, Phys. Rev. Lett. **72**, 1510 (1994).
 - ¹⁰ S. C. Zhang and D. P. Arovas, Phys. Rev. Lett. **72**, 1886 (1994).
 - ¹¹ A. G. Aronov, A. D. Mirlin, and P. Wölfle, Phys. Rev. B **49**, 16609 (1994).
 - ¹² D. Z. Liu, X. C. Xie, S. Das Sarma, and S. C. Zhang, Phys. Rev. B **52**, 5858 (1995).
 - ¹³ D. N. Sheng and Z. Y. Weng, Phys. Rev. Lett. **75**, 2388 (1995).
 - ¹⁴ Y. B. Kim, A. Furusaki and D. K. K. Lee, Phys. Rev. B **52**, 16646 (1995).
 - ¹⁵ J. Miller and J. Wang, Phys. Rev. Lett. **76**, 1461 (1996).
 - ¹⁶ B. Kramer, D. Belitz and M. Batsch, preprint (cond-

mat/9607043).

- ¹⁷ J. T. Chalker and P. D. Coddington, J. Phys. C **21**, 2665 (1988).
- ¹⁸ D. J. Thouless, M. Kohmoto, M. P. Nightingale and M. Den Nijs, Phys. Rev. Lett. **49**, 405 (1982).
- ¹⁹ Q. Niu, D. J. Thouless and Y. S. Wu, Phys. Rev. B **31**, 3372 (1985).
- ²⁰ D. P. Arovas, R. N. Bhatt, F. D. M. Haldane, P. B. Littlewood and R. Rammal, Phys. Rev. Lett. **60**, 619 (1988).
- ²¹ Y. Huo and R.N. Bhatt, Phys. Rev. Lett. **68**, 1375 (1992).
- ²² K. Yang and R.N. Bhatt, Phys. Rev. Lett. **76**, 1316 (1996).
- ²³ This relation is slightly different from the uniform field case,^{21,22} in which case one has $y = 1 - 1/(2\nu)$. This is because of the fact in that case there are only individual critical energies near which $\xi \sim |E - E_c|^{-\nu}$, for $w < w_c$.²² In the present case however, at $w = w_c$, the distance to criticality is proportional to E^2 , instead of E ($E_c = 0$), due to the fact that there is a *band* of extended states for $w < w_c$. Thus $\xi \sim |E - E_c|^{-2\nu}$ for $w = w_c$. For $w < w_c$ we still have $\xi \sim |E - E_c|^{-\nu}$, where E_c is the mobility edge. See Fig. 2.
- ²⁴ Muyu Guo and R. N. Bhatt, (in preparation).

FIG. 1. Number of extended states N_c versus system size N_s , for $h = 0.5$ and various w on a double logarithmic scale.

FIG. 2. Schematic phase diagram of the random flux problem

FIG. 3. The scaling functions $F^\pm(N_s|w - w_c|^{2\nu})$.

FIG. 4. Disorder averaged density of states $\rho(E)$ and density of current carrying states $\rho_c(E)$ at $w = 3.0$ and $h = 0.5$, for $L = 4$ and $L = 16$.











